



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF HELMINTHOSPORIUM, ANTHRACNOSE, PYTHIUM,
RHIZOCTONIA ROT AND SCLEROTINIA ROT IN NURSERY STOCK (NON-FOOD)**

PERMIT NUMBER – PER95488

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 30 MAY 2025 TO 31 MAY 2028

Permit Holder:

GREENLIFE INDUSTRY AUSTRALIA LIMITED
c/o AgAware Consulting Pty Ltd.
21 Rosella Avenue
STRATHFIELDSAYE VIC 3551

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

RAPIDOL TURF FUNGICIDE (APVMA No. 89251)

Containing: 250 g/L MANDESTROBIN as the only active constituent.

RESTRAINT:

DO NOT apply by aircraft.

DO NOT apply through any type of irrigation system or ultra-low volume spray system.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of runoff for at least 3 days after application.

DO NOT apply to vegetable or herb seedlings in the nursery.

SPRAY DRIFT RESTRAINTS:

Specific definitions for terms used in this section can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets not smaller than a MEDIUM spray droplet size category
- minimum distances between the application site and downwind sensitive areas (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones (metres)				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to maximum permit rate	0.5 m or lower	0	0	0	0	Not yet assessed
	1.0 m or lower	5	20	0	5	Not yet assessed

DO NOT apply by a vertical sprayer unless the following requirements are met:

- spray is not directed above the target canopy
- the outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site

Directions for Use:

Crop	Disease	Rate
NURSERY STOCK (NON-FOOD) - seedlings, tubes & plugs, potted colour, trees and shrubs, foliage plants, palms, grasses, fruiting plants (non-bearing) and ornamentals	Helminthosporium (<i>Bipolaris</i> spp.)	40 mL/100 m ²
	Anthracnose (<i>Colletotrichum</i> spp.)	(4 L/ha or 400 mL/100 L at 1000 L/ha)
	Pythium (<i>Pythium</i> spp.)	
	Rhizoctonia rot (<i>Rhizoctonia</i> spp.)	
	Sclerotinia rot (<i>Sclerotinia</i> spp.)	

Critical Use Comments:

- Rapidol Turf is more effective when used preventatively.
- If used as a preventative treatment, begin applications when conditions are favourable for disease development, prior to the onset of symptoms. If used as an early curative program, spray as soon as possible after the first symptoms are detected.
- Apply foliar spray via air-blast sprayer, air-shear sprayer, boom sprayer, track-sprayer or equivalent.
- Apply in sufficient water volume to achieve thorough coverage of all leaf surfaces and soil surfaces. Use a minimum of 400 L/ha. For maximum disease protection, ensure that all plant parts and soil surfaces are sufficiently covered by the spray solutions.
- DO NOT apply more than 2 applications of Rapidol Turf (or any other Group 11 fungicide) in any crop.
- Apply a second application at 14-21 days intervals, if required. DO NOT apply less than 14 days after the initial treatment.

Withholding Period:

DO NOT feed treated plants or prunings from any treated area to poultry or livestock.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the Agricultural and Veterinary Chemicals Code Act 1994, in accordance with the obligation imposed by that section.

To Avoid Crop Damage

The sensitivity of all nursery stock plants to be treated under this permit has not been fully evaluated. It is advisable to only treat a small area to ascertain the reaction before treating the whole crop.

Issued by the Australian Pesticides and Veterinary Medicines Authority