



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF ALTERNARIA BLIGHT, GREY MOULD, POWDERY
MILDEW AND SCLEROTINIA MOULD IN NURSERY STOCK (NON-FOOD)**

PERMIT NUMBER – PER95487

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 24 APRIL 2025 TO 30 APRIL 2028

Permit Holder:

GREENLIFE INDUSTRY AUSTRALIA LIMITED
c/o AgAware Consulting Pty Ltd
21 Rosella Avenue
STRATHFIELDSAYE VIC 3551

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

MIRAVIS PRIME ADEPIDYN TECHNOLOGY FUNGICIDE (APVMA No. 88887)

Containing: 250 g/L FLUDIOXONIL and 150 g/L PYDIFLUMETOFEN as the only active constituents.

RESTRAINT:

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of runoff for at least 3 days after application.

DO NOT use on vegetable or herb seedlings in the nursery.

Refer to spray drift restraints on product label.

Directions for Use:

Crop	Disease	Rate
NURSERY STOCK (NON-FOOD) seedlings, tubes & plugs, potted colour, trees and shrubs, foliage plants, palms, grasses, fruiting plants (non-bearing) and ornamentals	Alternaria blight (<i>Alternaria</i> spp.) Grey mould (<i>Botrytis</i> spp.) Powdery mildew (<i>Podosphaera</i> / <i>Erysiphe</i> spp.) Sclerotinia mould (<i>Sclerotinia</i> spp.)	750-1000 mL/ha or 75-100 mL/100 L at 1000 L/ha

Critical Use Comments:

- Use preventatively. Begin applications when conditions are favourable for disease development, prior to the onset of symptoms.
- Use the higher rate under conditions of high disease pressure and/or where a longer period of control is required.
- Apply foliar application via air-blast sprayer, air-shear sprayer, boom sprayer, track-sprayer or equivalent.
- Apply in sufficient water volume to achieve thorough coverage of all leaf surfaces.
- DO NOT apply more than 2 applications per crop.
- DO NOT apply less than 7 days after the initial treatment.
- DO NOT apply more than 2 applications of Miravis Prime (or any other Group 7 fungicide) in any crop.

Withholding Period:

DO NOT feed treated plants or prunings from any treated area to poultry or livestock.

Re-entry period:

Nursery stock: After application, do not enter treated areas for 3 days when performing high exposure activities, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

All other activities: After application, do not enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the Agricultural and Veterinary Chemicals Code Act 1994, in accordance with the obligation imposed by that section.

To Avoid Crop Damage

Nursery stock and cut flowers: This product has demonstrated toxicity under certain situations, application methods, application rates and weather conditions to nursery stocks including Impatiens, New Guinea Impatiens, Geraniums, Ferns, Anemone, Begonia, Kalanchoe, Saint Paulia and *Exacum affine*. This product may also leave undesirable spray residue on foliage and flowers. It is recommended that users treat a sample number of plants or a small area prior to whole crop treatment and monitor any phytotoxic effects that will compromise production goals. This action cannot guarantee crop safety as application, environmental and crop conditions may vary from test treatment to whole of crop treatment.

Issued by the Australian Pesticides and Veterinary Medicines Authority